

[Name of Document]      Specification

[Title of the Invention]

PHARMACEUTICAL COMPOSITION FOR USE IN  
TREATMENT OR PROPHYLAXIS OF ISCHEMIC HEART DISEASE

[Claims]

[Claim 1]    A pharmaceutical composition for use in treatment or prophylaxis of ischemic heart disease, the pharmaceutical composition containing a substance, as an active ingredient, which can increase intracellular cGMP production by acting on a natriuretic peptide receptor, and which has an effect of reducing an infarct region.

[Claim 2]    The pharmaceutical composition of claim 1, for suppressing ischemia-reperfusion injury in the treatment of ischemic heart disease.

[Claim 3]    The pharmaceutical composition of claim 1 or 2, wherein the ischemic heart disease is myocardial infarction.

[Claim 4]    The pharmaceutical composition of any one of claims 1 to 3, wherein the substance as the active ingredient is a natriuretic peptide or its salt.

[Claim 5]    The pharmaceutical composition of claim 4, wherein the natriuretic peptide is atrial natriuretic peptide.

[Detailed Description of the Invention]

[0001]

[Technical Field of the Invention]

This invention relates to a pharmaceutical composition for reducing an infarct region resulting from the ischemic necrosis of cells, the pharmaceutical

composition containing a substance, as an active ingredient, which can increase intracellular cGMP production by acting on a natriuretic peptide receptor.

[0002]

[Prior Art]

In recent years, ischemic heart disease has posed a major problem in an aging population. Of cardiac diseases which are diseases of circulatory organs, myocardial infarction ascribed to cardiovascular disorder, in particular, is a serious, potentially fatal disease which either obstructs the coronary artery or substantially decreases the blood flow resulting in ischemic necrosis of myocytes and deteriorating cardiac function. The direct cause of myocardial infarction is a decrease or interruption of the blood flow to the myocardium due to coronary arteriosclerosis or thrombus formation in the coronary artery. The disease can result in either acute or chronic cardiac failure.

[0003]

Methods adopted for treatment of ischemic heart disease include the dilatation of the obstructed coronary artery by use of an intravascularly inserted balloon, maintenance of blood flow by intravascular insertion of a stent, and dissolution and removal of a thrombus formed in the blood vessel with the use of a thrombolytic agent. With any of such treatments, it is known that as blood flow is restored in the coronary artery, Ca overload or free radicals occur, increasing the region of cellular necrosis.

Prevention of the occurrence of such ischemia-reperfusion injury is difficult, and no effective method of treatment has been established.

[0004]

[Problems to be Solved by the Invention]

The present invention provides a pharmaceutical composition for reducing an infarct region resulting from the ischemic necrosis of cells, the pharmaceutical composition containing a substance, as an active ingredient, which can increase intracellular cGMP production by acting on a natriuretic peptide receptor.

More specifically, the invention provides a pharmaceutical composition for suppressing ischemia-reperfusion injury in the treatment of ischemic disease.

[0005]

[Means for Solving the Problems]

hANP, a natriuretic peptide, is used as an agent of symptomatic therapy for alleviating symptoms of cardiac failure, because it has a diuretic action, and exhibits a blood pressure lowering effect by promoting production of cGMP, which is considered to be a second messenger of relaxation in vascular smooth muscle cells, to induce relaxation of blood vessels (e.g., coronary artery).

[0006]

The inventors of the present invention further studied the properties of natriuretic peptides, and found for the first time that these peptides can reduce an infarct region occurring in a model of acute myocardial

infarction involving ischemia reperfusion. This finding led them to accomplish this invention.

[0007]

That is, the present invention relates to a pharmaceutical composition for use in the treatment or prophylaxis of ischemic heart disease, such as myocardial infarction, the pharmaceutical composition containing a substance, as an active ingredient, which can increase intracellular cGMP production by acting on a natriuretic peptide receptor, and which has the effect of reducing an infarct region. In the present invention, "to reduce an infarct region" means to suppress enlargement of an infarct region.

[0008]

The present invention also relates to a method for treatment or prophylaxis of ischemic disease, comprising administering a substance to a patient with ischemic disease, which substance can increase intracellular cGMP production by acting on a natriuretic peptide receptor, and which has the effect of reducing an infarct region.

[0009]

Whether a certain substance can become the active ingredient of a pharmaceutical composition for use in the treatment or prophylaxis of ischemic disease, the pharmaceutical composition related to the present invention, can be investigated by using a known method, for example, the methods described in Minamitake, Y., et al., Biochem. Biophys. Res. Commun., 172, 971-978 (1990); Furuya, M., et

al., Biochem. Biophys. Res. Commun., 170, 201-208 (1990); Furuya, M., et al., Biochem. Biophys. Res. Commun., 177, 927-931 (1991); Hidaka, H. et al., Folia Pharmacologica Japonica, 101, 309-325 (1993).

[0010]

Preferred as the substance as an active ingredient according to the present invention are natriuretic peptides. Of them, natriuretic peptide (ANP) and brain natriuretic peptide (BNP) are preferred, and ANP is the most preferred.

[0011]

As ANP, there can be used human ANP (human atrial natriuretic peptide; hANP, Kangawa et al., Biochem. Biophys. Res. Commun., Vol. 118, p. 131, 1984) (Seq. ID No. 1) or rat ANP (Kangawa et al., Biochem. Biophys. Res. Commun., Vol. 121, p. 585, 1984) (Seq. ID No. 2), each ANP comprising 28 amino acids. The peptide as the active ingredient in the present invention may be a peptide having a ring structure of ANP (formation of a disulfide bond based on Cys), and a C-terminal portion succeeding the ring structure. An example of such a peptide is a peptide having amino acid residues at the 7-position to the 28-position of ANP (Seq. ID No. 3). Another example is frog ANP (Seq. ID No. 5). Of them, human ANP (hANP) is particularly preferred.

[0012]

An example of BNP is human BNP comprising 32 amino acids and involving the formation of a disulfide bond, like the above-described ANP (Sudoh et al., Biochem. Biophys.

Res. Commun., Vol. 159, p. 1420, 1989) (Seq. ID No. 4). Various BNP's of the origin other than human, such as pig BNP (Seq. ID No. 6) and rat BNP (Seq. ID No. 7), are also known, and can be used similarly. A further example is chicken BNP (Seq. ID No. 8).

[0013]

Examples of CNP are pig CNP comprising 22 amino acids and involving the formation of a disulfide bond, like the above-described ANP and BNP (Sudoh et al., Biochem. Biophys. Res. Commun., Vol. 168, p. 863, 1990) (Seq. ID No. 9; human and rat also have the same amino acid sequence), chicken CNP (Arimura et al., Biochem. Biophys. Res. Commun., Vol. 174, p. 142, 1991) (Seq. ID No. 10), and frog CNP (Yoshihara et al., Biochem. Biophys. Res. Commun., Vol. 173, p. 591, 1990) (Seq. ID No. 11).

[0014]

Furthermore, any person skilled in the art can apply modification, such as deletion, substitution, addition or insertion, and/or chemical modification to amino acid residues in the amino acid sequence of a known natriuretic peptide (e.g., the aforementioned human ANP; hANP), as desired, by a known method. One skilled in the art can confirm that the resulting compound is a compound which has the activity of acting on a receptor of ANP to increase cGMP production. Derivatives having this activity, therefore, are included in the substance as an active ingredient in accordance with the present invention. Moreover, the substances involved in the present invention

are not restricted to the above peptides, as long as they are substances capable of acting on a natriuretic peptide receptor to increase intracellular cGMP production. These substances may be non-peptide compounds.

[0015]

The substance as an active ingredient according to the present invention may be of a free type, or its pharmaceutically acceptable salt. The salt with an inorganic acid includes, for example, salts with hydrochloric acid, sulfuric acid, and phosphoric acid. The salt with an organic acid may, for example, be acid addition salts with formic acid, acetic acid, butyric acid, succinic acid, and citric acid. The salt may be in the form of a metal salt with sodium, potassium, lithium or calcium, or a salt with an organic base.

[0016]

The substance as an active ingredient is preferably mixed with known pharmaceutically acceptable carriers, vehicles, or diluents, and administered by an administration method used generally for drugs, for example, an oral administration method, or a parenteral administration method, such as intravenous administration, intracoronary administration, intramuscular administration, or subcutaneous administration. The pharmaceutical composition of the present invention can be produced, for example, by mixing, as desired, the active ingredient, pharmaceutically acceptable carriers, flavors, vehicles, and stabilizers. To produce solid preparations for oral administration, such

as tablets, capsules, granules, and fine granules, the following additives can be used: (1) vehicles such as lactose, starch, and microcrystalline cellulose, (2) binders such as hydroxypropylcellulose, and polyvinylpyrrolidone, (3) disintegrants such as starch and crosscarmellose sodium, (4) plasticizers such as macrogol and triethyl citrate, (5) lubricants such as magnesium stearate and talc, (6) coating materials such as hydroxypropyl methylcellulose, and Eudragit, and (7) taste correctives such as sucrose and mannitol, odor correctives, and colorants.

[0017]

To produce injections, ophthalmic solutions, or transnasal preparations, the following additives can be added: (1) tonicity agents such as sodium chloride, D-mannitol, and D-sorbitol, (2) pH regulators such as hydrochloric acid and citric acid, (3) buffering agents such as sodium citrate, sodium acetate, and boric acid, and (4) soothing agents such as procaine hydrochloride; as well as stabilizers, and surface active agents. In consideration of the stability, etc. of the active ingredient, it can be selected whether the active ingredient should be formed into a preparation to be used after dissolution or suspension when required, or into a liquid preparation.

[0018]

To produce preparations for external use, such as ointments and cataplasms, the following materials can be

added: (1) bases such as liquid petrolatum, petrolatum, and hydrophilic ointments, (2) emulsifying agents such as polysorbate 80, and tragacanth, (3) preservatives such as sodium benzoate, and propyl p-hydroxybenzoate, and (4) soothing agents such as procaine hydrochloride, stabilizers, and surface active agents.

[0019]

When the substance as an active ingredient is a natriuretic peptide, this peptide orally administered is degraded in the digestive tract, and thus this mode of administration is generally not effective. However, the peptide can be orally administered in the form of a preparation minimally degraded in the digestive tract, for example, microcapsules comprising the peptide, as the active ingredient, enclosed in a liposome. A mode of administration by absorption through the mucosa other than the digestive tract, such as the rectum or a sublingual area, is also possible. In this case, a dosage form, such as a suppository or a sublingual tablet, can be used for administration.

[0020]

The dose of the pharmaceutical composition of the present invention differs according to the age, the body weight, the severity of symptoms of, and the route of administration in, a patient with myocardial infarction or a patient potentially developing myocardial infarction. When the substance as an active ingredient is a natriuretic peptide, the pharmaceutical composition can be administered

at a dose of 0.01  $\mu\text{g/kg/min}$  to 0.2  $\mu\text{g/kg/min}$ , and is preferably administered in a dose of 0.025  $\mu\text{g/kg/min}$  to 0.1  $\mu\text{g/kg/min}$ , by the continuous intravenous route.

[0021]

[Example]

The following example shows that hANP, a natriuretic peptide, reduces the region of myocardial infarction occurring in models of acute myocardial infarction involving ischemia reperfusion.

#### Method

Thoracotomy was performed in 12 adult beagles weighing 14 to 23 kg under anesthesia with pentobarbital sodium, and hANP (0.1  $\mu\text{g/kg/min}$ ) was continuously administered for 10 minutes into the left anterior descending branch (LAD) of the coronary artery. Then, the LAD was completely obstructed into an ischemic state until the LAD was reperfused 90 minutes later. hANP (0.1  $\mu\text{g/kg/min}$ ) was continuously administered into the LAD over the course of 1 hour since 10 minutes before initiation of reperfusion. After 6 hours of reperfusion, a region at risk of developing infarction was evaluated by Evans blue staining, and the region of infarction was evaluated by TTC staining (Group A: 5 dogs). After 80 minutes of ischemia, the amount of endocardial collateral blood flow was measured by the microsphere method.

[0022]

A group receiving physiological saline, instead of hANP, into the LAD was provided as a control group (Group

B: 7 dogs).

[0023]

A protocol for the experiments is shown in FIG. 1. In the drawing, CPP denotes coronary perfusion pressure. The protocol shows that blood flowed in the LAD before start of the test, then the blood flow was interrupted for 90 minutes, then blood flow was restored again, and the test was completed at 360 minutes.

[0024]

The amount of endocardial collateral blood flow during ischemia, the mean blood pressure, and the heart rate were also measured to investigate whether or not these parameters took part in the effect of the hANP according to the present invention.

### Results

(1) FIG. 2 shows the size of the region at risk of myocardial infarction in the left ventricle of each test group. There was no significant difference between the test groups in the size of the region at risk of myocardial infarction.

[0025]

(2) In the models of acute myocardial infarction involving ischemia reperfusion, the hANP administration reduced the region of myocardial infarction. As shown in FIG. 3, the region of myocardial infarction in the control group (Group B) was  $41 \pm 3\%$  of the region at risk of myocardial infarction, while the region of myocardial infarction significantly decreased to  $21 \pm 5\%$  in the hANP

group (Group A)(FIG. 3).

[0026]

(3) No difference was confirmed between the groups in terms of the amount of endocardial collateral blood flow during ischemia. Moreover, changes in the heart rate and the mean blood pressure in each group were measured 5 and 10 minutes after start of the test, 90 minutes after initiation of ischemia, and 360 minutes after start of reperfusion. The mean blood pressure and the heart rate were confirmed to remain unchanged following administration of hANP.

[0027]

The above findings demonstrate that the administration of a natriuretic peptide suppresses ischemia-reperfusion injury in the treatment of ischemic disease. Thus, the effect of reducing the region of myocardial infarction is confirmed to be ascribed to the natriuretic peptide's action of reducing the region of myocardial infarction.

[0028]

[Sequence Listing]

#### SEQUENCE LISTING

<110> SUNTORY LIMITED

<120> Pharmaceutical Composition for Use in Treatment or  
Prophylaxis of Ischemic Heart Disease

<130> 000687

<160> 11